arXiv:math/0605468v1 [math.DG] 17 May 2006

A closed symplectic four-manifold has almost Kähler metrics of negative scalar curvature

Jongsu Kim [*]
Department of Mathematics, Sogang University
Seoul, 121-742, KOREA

February 2, 2008

Abstract

We show that every closed symplectic four-dimensional manifold (M, ω) admits an almost Kähler metric of negative scalar curvature compatible with ω.

1 Introduction

In Riemannian geometry an interesting problem related to a curvature is whether every smooth manifold of dimension bigger than 2 admits a metric with the curvature negative. Lohkamp's beautiful argument in [12] resolved this for the Ricci curvature case, while the relatively simpler case of scalar curvature had been understood earlier, see [2, Chap. 4].

Meanwhile, the extensive study on symplectic manifolds in the last decades has broadened the knowledge on the field and supplied large families of closed symplectic manifolds [6, 7]. Therefore the study of Riemannian metrics compatible to a symplectic structure, so-called almost Kähler metrics, re-emerges as an interesting and promising subject. Remarkably, the study of almost

---

[*] Supported by grant No. R14-2002-044-01000-0(2002) from the Korea Science and Engineering Foundation. Keywords: Almost Kähler metric, symplectic structure, scalar curvature. MS Classification(2000): 53C15, 53C20, 53D35

Kähler metrics in Riemannian geometry has a long history, from the works of 1960's or earlier [14] to recent works in several directions [1], [8], [10], etc..

Motivated by this, in this paper we study the existence of negative scalar-curved almost Kähler metrics on a symplectic manifold. In the general Riemannian metric case afore-mentioned, the main argument dealing with scalar curvature was to use a conformal deformation. For almost Kähler metrics the latter can not be used because it does not preserve the symplectic structure. So even the scalar curvature question does not seem trivial in almost Kähler case.

Our argument is based on the Lohkamp's framework in [12] and in its course there were two main ingredients to proving the following main theorem in this paper;

Theorem 1 *Every closed symplectic four-dimensional manifold* (M, ω) *admits an* ω*-compatible almost Kähler metric of negative scalar curvature.*

The first ingredient is to find an almost Kähler *island* metric, i.e. an almost Kähler metric on $\mathbb{R}^n$ which has the scalar curvature negative on a pre-compact open subset and is Euclidean outside of its closure.

The second one is to find an almost Kähler deformation to decrease the scalar curvature outside copies of the island metrics which are embedded into the (enlarged) almost Kähler manifold by a metric surgery. This deformation corresponds to the conformal deformation [12, p. 669] of Lohkamp's.

As our almost Kähler deformation completely avoids any conformal deformation, it brings about new technical problems. One remarkable difference from Lohkamp's argument is that all the metrics involved, including the island metrics, should be locally close to Euclidean metrics. It is because we resorted to specific local orthonormal frame fields $\{\omega_i\}$ in defining the deformation, and these fields are manageable if the metrics are close to Euclidean ones. This technical aspect is the reason why we restricted to real 4-dimension in this article, though we believe the higher dimensional case should be provable by some modification.

One may view the sort of result like Theorem 1 as indicating that the space of almost Kähler metrics on a symplectic manifold may be big enough to contain rich geometric features. And one may speculate on any implication of almost Kähler geometry to the topology of symplectic manifolds.

The paper is organized as follows. In section 2 we provide preliminaries for almost Kähler metrics, an estimate for injectivity radius and the Besicovitch covering. In section 3 we describe the local orthonormal frame fields $\{\omega_i\}$ of almost Kähler metrics which are close to Euclidean metrics. In section 4 we explain the almost Kähler deformation depending on $\{\omega_i\}$ which decreases the scalar curvature away from a compact set. In section 5 we describe an *island* metric and modify it with the result of section 4. In section 6 we do a surgery of embedding many copies of metrics of section 5 into the given almost Kähler manifold. In section 7 we apply the deformation of the section 4 to the resulting manifold of section 6 and prove the main theorem. There is Appendix which proves some estimates on $\{\omega_i\}$ and derives a scalar curvature formula.

Acknowledgement. The author would like to thank Yutae Kang for many helpful discussions on this work and thank the referee for his suggestions which made this paper more readable.

2 Preliminaries

In this section we explain definitions and formulas which will be needed in later sections.

An *almost-Kähler* metric is a Riemannian metric g compatible with a symplectic structure ω on a smooth manifold, i.e. $\omega(x, y) = g(Jx, y)$ for an almost complex structure J, where x, y are tangent vectors at a point of the manifold. We often denote it by the triple (g, ω, J) or the couple (g, ω) to specify what ω or J is. Notice that any one of the pairs (g, ω), (ω, J), or (g, J) determines the other two. For a fixed ω, we shall call a metric g ω-almost Kähler if g is compatible with ω and denote by Ω_ω the set of all smooth ω-almost Kähler metrics. An almost-Kähler metric (g, ω, J) is Kähler if and only if J is integrable.

An almost complex structure J gives rise to a type decomposition of any symmetric (2,0)-tensor h which can be written as $h = h^+ + h^-$, where h^+ and h^- are symmetric 2-tensors defined by $h^+(x, y) = \frac{1}{2}\{h(x, y) + h(Jx, Jy)\}$ and $h^-(x, y) = \frac{1}{2}\{h(x, y) - h(Jx, Jy)\}$ for two vectors x and y tangent to M. A symmetric $(2, 0)$-tensor h is called *J-invariant* or *J-anti-invariant* if $h = h^+$ or $h = h^-$ respectively.

The set Ω_ω is naturally an infinite dimensional Fréchet manifold. For a smooth curve g_t in Ω_ω with the corresponding curve J_t of almost complex structures, $h = \frac{dg_t}{dt}|_{t=0}$ is J_0-anti-invariant. Conversely, for g in Ω_ω with the corresponding J, any J-anti-invariant symmetric $(2,0)$-tensor field h is tangent to a smooth curve in Ω_ω. More precisely, $t \mapsto ge^{ht}$ is such a smooth curve in Ω_ω, where ge^{ht} is defined by $ge^{ht}(x,y) = g(x, e^{(g^{-1}h)t}y)$, i.e.

$$ge^{ht}(x,y) = g(x,y) + g(x, \sum_{k=1}^{\infty} \frac{t^k (g^{-1}h)^k (y)}{k!}), \tag{1}$$

where $g^{-1}h$ is the lifted $(1,1)$-tensor of h with respect to g. Given $g \in \Omega_\omega$ with corresponding J, any other metric $\tilde{g}$ in Ω_ω can be expressed as

$$\tilde{g} = ge^h, \tag{2}$$

where h is a J-anti-invariant symmetric $(2,0)$-tensor field uniquely determined. See [3] for details.

We recall local lower bound estimates on the injectivity radius of a manifold. For p in a complete Riemannian manifold (N,g), let l_p denote the length of the shortest geodesic loop based at p. If the sectional curvature $K_N \leq \Lambda^2$ on $B^g_r(p)$, the open geodesic ball of radius r centered at p w.r.t. g, and $r \leq \frac{1}{2} l_p$, then the injectivity radius $\mathrm{inj_g(p)}$ of (N,g) at p satisfies $\mathrm{inj_g(p)} \geq \mathrm{min(r, \frac{\pi}{\Lambda})}$ by an estimate of Klingenberg [11, Lemma 1] and the fact that if a point q is conjugate to p then their distance $d(p,q) \geq \frac{\pi}{\Lambda}$, [4, p. 30].
Now suppose $K_N \leq \Lambda^2$ on $B^g_r(p)$, where $r \leq \frac{\pi}{\Lambda}$. Choose any numbers r_0 and s such that $r_0 + 2s \leq r$, $r_0 \leq \frac{r}{4}$. Then one has the estimate due to Cheeger-Gromov-Taylor [5, Theorem 4.3]

$$l_p \geq r_0 [1 + \frac{v^{-\Lambda}_{r_0+s}}{v^g_s(p)}]^{-1}. \tag{3}$$

where $v^g_s(p)$ is the g-volume of $B^g_s(p)$ and $v^{-\Lambda}_{r_0+s}$ is the volume of the geodesic ball of radius $r_0 + s$ in the space form of constant curvature $-\Lambda$.

Next we recall the Besicovitch covering [12, Prop 4.4];

Lemma 1 *Let (M,h) be a Riemannian manifold with the injectivity radius* $\mathrm{inj_h} \geq 100$ *and the sectional curvature $|K_h| < 1$. Then there are constants $\kappa \in Z^{>0}$ and $m_0 \geq 1$ depending only on the dimension of M such that*

(i) For each $m \geq m_0$ there is a set $A = A(m) \subset M$ with $d_{m^2h}(a,b) > 5$ for $a \neq b, a, b \in A(m)$.

(ii) $\mathcal{A}(m) = \{\overline{B_5^{m^2h}(a)} \mid a \in A(m)\}$ is a (closed) covering of M and splits into κ disjoint families $\mathcal{B}_j$ with $\overline{B_{10}^{m^2h}(a)} \cap \overline{B_{10}^{m^2h}(b)} = \phi$ if $\overline{B_5^{m^2h}(a)}$ and $\overline{B_5^{m^2h}(b)}$ belong to the same $\mathcal{B}_j$.

We shall use the following notation. When h is a tensor field on a domain V, we denote the C^k norm of h with respect to a metric g on V by $\|h\|_{C_g^k(V)}$. Or sometimes we express it by writing "$\|h\|_{C_g^k}$ on V". Here $\|h\|_{C_g^k(V)} = \sum_{i=0}^k \|(\nabla_g)^i h\|_{C_g^0(V)}$ and ∇_g is the Levi-Civita connection of g.

3 Almost-Kähler metrics which are close to Euclidean metrics

We start with a Euclidean metric g_0 on $\mathbb{R}^4$ or an open subset Ω. We assume that there exists a point p in Ω and that $B_{10^4}^{g_0}(p)$, the open g_0-ball centered at p with radius 10^4, is in Ω. Now suppose that there is another Riemannian metric g on Ω and $\|g - g_0\|_{C_{g_0}^0(B_{10^4}^{g_0}(p))} \leq \varepsilon$, where ε is a small positive number, say $\varepsilon < 10^{-4}$.

For another point $q \in B_{10^4}^{g_0}(p)$ and a minimal geodesic $\gamma_{g_0}(t)$ of g_0 connecting $\gamma_{g_0}(0) = p$, $\gamma_{g_0}(t_0) = q$, we have

$$d_{g_0}(p,q) = \int_0^{t_0} |\gamma_{g_0}'(s)|_{g_0} ds \geq \int_0^{t_0} \frac{|\gamma_{g_0}'(s)|_g}{\sqrt{1+\varepsilon}} ds \geq \frac{d_g(p,q)}{\sqrt{1+\varepsilon}},$$

where $d_g(p,\cdot)$ and $d_{g_0}(p,\cdot)$ are the distance functions from p induced by g and g_0, respectively. To prove the inequality in the opposite direction, we first show $B_{9999}^g(p) \subset B_{10^4}^{g_0}(p)$. Let's consider a g-geodesic $\gamma(t)$ with $\gamma(0) = p$ and $|\gamma'(t)|_g \equiv 1$. For a small number δ with $0 < \delta \ll \varepsilon$, let $t_1 > 0$ be the first positive value of t such that $\gamma(t)$ is on the boundary of $B_{10^4-\delta}^{g_0}(p)$. Then

$$10^4 - \delta = d_{g_0}(p, \gamma(t_1)) \leq \int_0^{t_1} |\gamma'(s)|_{g_0} ds \leq \frac{1}{\sqrt{1-\varepsilon}} \int_0^{t_1} |\gamma'(s)|_g ds = \frac{1}{\sqrt{1-\varepsilon}} t_1.$$

As $9999 < (10^4 - \delta)\sqrt{1-\varepsilon} \leq t_1$, this implies $B_{9999}^g(p) \subset B_{10^4}^{g_0}(p)$. Now for $q \in B_{9999}^g(p)$ and a minimal geodesic $\gamma_g(t)$ of g connecting $\gamma_g(0) = p$, $\gamma_g(t_2) = q$, we have

$$d_g(p,q) = \int_0^{t_2} |\gamma_g'(s)|_g ds \geq \int_0^{t_2} \sqrt{1-\varepsilon} \cdot |\gamma_g'(s)|_{g_0} ds \geq \sqrt{1-\varepsilon} \cdot d_{g_0}(p,q).$$

In sum we proved;

Lemma 2 *Suppose that a Euclidean metric* g_0 *and a Riemannian metric* g *on* $B^{g_0}_{10^4}(p) \subset \mathbb{R}^4$ *satisfies* $\|g - g_0\|_{C^0_{g_0}(B^{g_0}_{10^4}(p))} \leq \varepsilon < 10^{-4}$*. Then,* $B^{g}_{9999}(p) \subset B^{g_0}_{10^4}(p)$ *and for any* $q \in B^{g}_{9999}(p)$*, we have*

$$\sqrt{1-\varepsilon} \cdot d_{g_0}(p,q) \leq d_g(p,q) \leq \sqrt{1+\varepsilon} \cdot d_{g_0}(p,q). \tag{4}$$

One can prove the following lemma from (3) and (4). We omit the detail of the proof which is straightforward.

Lemma 3 *There exists* $\lambda_1 < 10^{-4}$ *such that if* $\|g - g_0\|_{C^2_{g_0}(B^{g_0}_{10^4}(p))} \leq \lambda_1$*, then* g *has the injectivity radius at* p*,* $\mathrm{inj_g(p)} \geq 200$*.*

Consider the symplectic structure $\omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_4$ on $\mathbb{R}^4 = \{(x_1, x_2, x_3, x_4) |\ x_i \in \mathbb{R}\}$ together with the compatible Euclidean metric $g_0 = \sum_{i=1}^4 dx_i \otimes dx_i$ and the (almost) complex structure J_0. We shall deform an ω-almost Kähler metric g on Ω containing $B^{g_0}_{10^4}(p)$, satisfying $\|g - g_0\|_{C^l_{g_0}(B^{g_0}_{10^4}(p))} \leq \varepsilon < 10^{-4}$, where $l \geq 4$ and ε is a positive number to be determined later. The Euclidean metric $(\mathbb{R}^4, g_0, \omega, J_0)$ has an orthonormal co-frame field $\omega^0_1 = dr_{g_0}$, $\omega^0_2 = J_0(dr_{g_0}) = r_{g_0}\sigma_1$, $\omega^0_3 = r_{g_0}\sigma_2$, $\omega^0_4 = r_{g_0}\sigma_3 = J_0(\omega^0_3)$, where $r_{g_0}(\cdot) = d_{g_0}(p, \cdot)$ and $\sigma_1, \sigma_2, \sigma_3$ is the canonical orthonormal co-frame field of the standard metric on S^3 which satisfy $d\sigma_1 = 2\sigma_2 \wedge \sigma_3, d\sigma_2 = 2\sigma_3 \wedge \sigma_1$ and $d\sigma_3 = 2\sigma_1 \wedge \sigma_2$. We shall find an orthonormal co-frame field of g near p: ω_i, $i = 1, 2, 3, 4$ which are close to ω^0_i. On a ball $B^g_{100}(p) - \{p\} \subset B^{g_0}_{101}(p)$ we define formally

$$\begin{aligned} &\omega_1 = dr_g, \qquad \omega_2 = J_g(dr_g), \\ &\omega_3 = \frac{\omega^0_3 - g(\omega^0_3, \omega_1)\omega_1 - g(\omega^0_3, \omega_2)\omega_2}{|\omega^0_3 - g(\omega^0_3, \omega_1)\omega_1 - g(\omega^0_3, \omega_2)\omega_2|_g}, \qquad \omega_4 = J_g(\omega_3). \end{aligned} \tag{5}$$

where $r_g(\cdot) = d_g(p, \cdot)$ and J_g is the almost complex structure determined by ω and g. We need the following proposition on some properties of $\{\omega_i\}$ whose proof is postponed to the Appendix;

Proposition 1 *There exist* λ_2 *with* $0 < \lambda_2 \leq \lambda_1$ *and a positive constant* C *such that if* $\|g - g_0\|_{C^l_{g_0}} \leq \lambda \leq \lambda_2$ *on* $B^{g_0}_{10^4}(p) \subset \mathbb{R}^4$*, where* l *is an integer* $4 \leq l \leq 2\kappa + 4$ *and* κ *is the constant (in dimension four) of the Besicovitch* *covering Lemma 1, then* $\omega_1, \omega_2, \omega_3, \omega_4$ *in the formula (5) are well defined and satisfy on* $B^g_{100}(p) - \{p\}$*;*

$$\|(\nabla^{g_0})^j \omega_i - (\nabla^{g_0})^j \omega_i^0\|_{C^0_{g_0}} < \frac{C \cdot \lambda}{r^j_{g_0}}, \qquad \textit{for } j = 0, 1, \cdots, l-2. \tag{6}$$

where ∇^{g_0} *is the Levi-Civita connection of* g_0*.*

For the above metric $g = \sum_{i=1}^4 \omega_i \otimes \omega_i$ on $B^g_{100}(p)$, consider the connection 1-forms ω_{ij} w.r.t. the co-frame ω_i: $d\omega_i = \sum_{j=1}^4 \omega_{ij} \wedge \omega_j$, with $\omega_{ij} = -\omega_{ji}$. Writing ω_{ij} as $\omega_{ij} = \sum_{k=1}^4 a_{ijk} \cdot \omega_k$, we get $d\omega_i = \sum_{j<k}(a_{ikj} - a_{ijk})\omega_j \wedge \omega_k$. So we get $2a_{ijk} = \langle d\omega_k, \omega_i \wedge \omega_j\rangle_g - \langle d\omega_i, \omega_j \wedge \omega_k\rangle_g - \langle d\omega_j, \omega_k \wedge \omega_i\rangle_g$, where $\langle \cdot, \cdot \rangle_g$ is the naturally induced metric on the space of 2-forms from g.

For the Euclidean metric $(\mathbb{R}^4, g_0, \omega, J_0)$ with the above co-frame $\{\omega_i^0\}$, the a_{ijk}'s are all zero except

$$\begin{cases} a_{122} & = a_{133} = a_{144} = -a_{212} = -a_{313} = -a_{414} = \frac{1}{r_{g_0}}, \\ a_{234} & = a_{342} = a_{423} = -a_{324} = -a_{432} = -a_{243} = -\frac{1}{r_{g_0}}. \end{cases} \tag{7}$$

For later we need the following estimate.

Lemma 4 *There exists* λ_3 *with* $\lambda_3 \leq \lambda_2$ *such that if* $\|g - g_0\|_{C^4_{g_0}} \leq \lambda_3$ *on* $B^{g_0}_{10^4}(p) \subset \mathbb{R}^4$*, then for* $i, j, k, l = 1, 2, 3, 4$*, the* a_{ijk} *of the* ω_i*'s in the formula* (5) *satisfy:*

$$|a_{ijk}| < \frac{2}{r_g}, \; |a_{ijk,l}| < \frac{2}{r_g^2} \textit{ on } B^g_{100}(p) - \{p\},$$

where $a_{ijk,l}$ is defined in the formula $d(a_{ijk}) = \sum_{l=1}^4 a_{ijk,l}\omega_l$.

Proof: One needs a straightforward computation: show $|\langle d\omega_i, \omega_j \wedge \omega_k\rangle_g - \langle d\omega_i^0, \omega_j^0 \wedge \omega_k^0\rangle_g| \leq \frac{\text{constant}\cdot\lambda}{r_{g_0}}$ from (6). Then $a_{ijk} = \frac{1}{2}(\langle d\omega_k, \omega_i \wedge \omega_j\rangle_g - \langle d\omega_i, \omega_j \wedge \omega_k\rangle_g - \langle d\omega_j, \omega_k \wedge \omega_i\rangle_g)$ satisfies $|a_{ijk} - a^0_{ijk}| \leq \frac{\text{constant}\cdot\lambda}{r_{g_0}}$. From (7), $|a_{ijk}| < \frac{2}{r_g}$ follows. One can do similarly for $|a_{ijk,l}|$.

4 Scalar curvature diffusion

We shall perturb locally an almost Kähler metric g which satisfies the hypothesis of Lemma 4 and is written as $g = \sum_{i=1}^4 \omega_i \otimes \omega_i$ on $B^g_{100}(p) - \{p\}$, where ω_i's are as defined in (5).

We use the functions of [12]; $f_{d,s}(t) \in C^\infty(\mathbb{R}, \mathbb{R}^{\geq 0})$ for $d, s > 0$ by $f_{d,s} = s \cdot \exp(-\frac{\mathrm{d}}{\mathrm{t}})$ on $\mathbb{R}^{>0}$ and $f_{d,s} = 0$ on $\mathbb{R}^{\leq 0}$. Also choose an $h \in C^\infty(\mathbb{R}, [0,1])$ with $h = 0$ on $\mathbb{R}^{\geq 1}$, $h = 1$ on $\mathbb{R}^{\leq 0}$ and $h^b_\epsilon(t) = h(\frac{1}{\epsilon}(t-b))$, $b > 0$, $\epsilon > 0$. Now we define a perturbed metric for $b + \epsilon < c \leq 9$;

$$\begin{aligned} g^{b,\epsilon,c}_{d,s} = & \frac{\omega_1 \otimes \omega_1}{\{1 + h^b_\epsilon(c - r_g) \cdot f_{d,s}(c - r_g)\}} + \{1 + h^b_\epsilon(c - r_g) \cdot f_{d,s}(c - r_g)\}\omega_2 \otimes \omega_2 \\ & + \omega_3 \otimes \omega_3 + \omega_4 \otimes \omega_4. \end{aligned} \tag{8}$$

Then $g^{b,\epsilon,c}_{d,s}$ is a smooth ω-almost Kähler metric on $B^{g_0}_{10^4}(p)$. We get

Lemma 5 *Suppose that an ω-almost Kähler metric g on a domain $\Omega \subset \mathbb{R}^4$ satisfies $\|g - g_0\|_{C^l_{g_0}(B^{g_0}_{10^4}(p))} \leq \lambda_3$ where $B^{g_0}_{10^4}(p) \subset \Omega$ and l is an integer $4 \leq l \leq 2\kappa + 4$. For each $a, b, \epsilon, c > 0$ such that $9 \geq c > b + \epsilon > b > a > 0$, consider the orthonormal co-frame ω_i as defined in* (5) and define $g^{b,\epsilon,c}_{d,s}$ on $B^{g_0}_{10^4}(p)$ as in (8). Then the following statements hold;
(i) For each $D > 0$, there is a constant $\alpha = \alpha(b, \epsilon, c, D)$ such that for $d \geq D$ and $s \in (0,1]$, $\|g^{b,\epsilon,c}_{d,s} - g\|_{C^{l-2}_{g_0}(B^{g_0}_{10^4}(p))} < s \cdot \alpha$,
(ii) For each $\varepsilon > 0$ there exists a $d = d(\varepsilon, c)$ such that for every $d \geq d(\varepsilon, c)$ and each $s \in (0,1]$, $\|g^{b,\epsilon,c}_{d,s} - g\|_{C^{l-2}_{g_0}(B^{g_0}_{10^4}(p))} < \varepsilon$ holds.

Proof. We apply the estimates (6) to

$$g^{b,\epsilon,c}_{d,s} - g = -\frac{h^b_\epsilon(c - r_g) \cdot f_{d,s}(c - r_g)}{1 + h^b_\epsilon(c - r_g) \cdot f_{d,s}} \omega_1 \otimes \omega_1 + h^b_\epsilon(c - r_g) \cdot f_{d,s}(c - r_g) \omega_2 \otimes \omega_2$$

which has support in $c - b - \epsilon \leq r_g \leq c$. The computation is elementary and we omit it. Here one may use $\|(\nabla^{g_0})^j dr_{g_0}\|_{C^0_{g_0}} < \frac{\text{constant}}{r^j_{g_0}}$, for $0 \leq j \leq l - 2$; see the formula (35) in the Appendix. ■

Note that λ_3 is a constant already determined, so in the statement of Lemma 5 we do not specify the dependence of α or d on λ_3.

For the metric $g^{b,\epsilon,c}_{d,s}$, we compute its scalar curvature using the orthonormal co-frame field $\tilde{\omega_1} = \frac{\omega_1}{\sqrt{1 + h^b_\epsilon \cdot f_{d,s}(c - r_g)}}$, $\tilde{\omega_2} = \sqrt{1 + h^b_\epsilon \cdot f_{d,s}(c - r_g)} \cdot \omega_2$, $\tilde{\omega_3} = \omega_3$ and $\tilde{\omega_4} = \omega_4$ and the formula $d\tilde{\omega}_i = \sum_{j=1}^4 \tilde{\omega}_{ij} \wedge \tilde{\omega}_j$.

Setting $Y = Y(r_g) = h^b_\epsilon(c - r_g) \cdot f_{d,s}(c - r_g)$, the scalar curvature of $g^{b,\epsilon,c}_{d,s}$ is computed as follows, see Appendix for details:

$$\begin{aligned} s(g^{b,\epsilon,c}_{d,s}) =& - Y'' - Y' \cdot (3a_{122} + 2a_{133} + 2a_{144}) - 2(Y+1)\{\sum_{i=2}^{4}(a_{1ii,1} + a^2_{1ii}) \\ & + a^2_{134} + a_{122}a_{133} + a_{122}a_{144} + a_{133}a_{144} + \frac{1}{4}(a_{243} - a_{234})^2\} \\ & - \frac{2}{Y+1}\{a_{233,2} + a_{244,2} + a^2_{233} + a^2_{244} + a_{233}a_{244} + \frac{1}{4}(a_{243} + a_{234})^2\} \\ & - \frac{1}{2}\sum_{i=3}^{4}\{Ya_{2i1} - (Y+2)a_{21i}\}^2 + I_0, \end{aligned} \tag{9}$$

where
$$\begin{aligned} \frac{I_0}{2} =& a_{232,3} + a_{242,4} + a_{343,4} + a_{434,3} - a^2_{232} - a^2_{242} - a^2_{343} - a^2_{434} + a_{242}a_{433} \\ & + a_{232}a_{344} + a_{342}a_{234} - a_{342}a_{243} + \frac{1}{2}a^2_{234} + \frac{1}{2}a^2_{243} - a_{243}a_{234}. \end{aligned}$$

The scalar curvature of g can be obtained by putting $Y = 0$ into (9). We are concerned with their difference:

$$s(g^{b,\epsilon,c}_{d,s}) - s(g) = -Y'' - Y'(3a_{122} + 2a_{133} + 2a_{144}) - Y \cdot I_1 \tag{10}$$

where

$$\begin{aligned} \frac{I_1}{2} =& \frac{1}{4}(a_{243} - a_{234})^2 + \sum_{i=2}^{4}(a_{1ii,1} + a^2_{1ii}) + a^2_{134} + a_{122}a_{133} + a_{122}a_{144} + a_{133}a_{144} \\ & + \frac{1}{4}\{Y(a^2_{231} + a^2_{241}) - 2(Y+2)(a_{213}a_{231} + a_{214}a_{241}) + (Y+4)(a^2_{213} + a^2_{214})\} \\ & - (\frac{1}{Y+1})\{a_{233,2} + a_{244,2} + a^2_{233} + a^2_{244} + a_{233}a_{244} + \frac{1}{4}(a_{243} + a_{234})^2\}. \end{aligned}$$

Due to the estimate of Lemma 4, we can prove that the above almost Kähler deformation diffuses the scalar curvature, analogously to [12, Proposition 2.1]

Lemma 6 *Suppose that an almost Kähler metric g on a domain Ω in $\mathbb{R}^4$ satisfies $\|g - g_0\|_{C^4_{g_0}(B^{g_0}_{10^4}(p))} \leq \lambda_3 < 1$. For each $a, b, \epsilon, c > 0$ such that*

$9 \geq c > b+\epsilon > b > a > 0$, *consider the orthonormal co-frame* ω_i *as in (5) and define* $g_{d,s}^{b,\epsilon,c}$ *on* $B_{10^4}^{g_0}(p)$ *as in (8). There are constants* $\gamma = \gamma(a,b,c) > 0$ *such that for* $d \geq \gamma$, $s \in [0,1]$ *the following statements hold:*

(i) $g_{d,s}^{b,\epsilon,c} \equiv g$ on $B_{c-b-\epsilon}^{g}(p)$,

(ii) $s(g_{d,s}^{b,\epsilon,c}) - s(g) \leq 0$ on $B_{10^4}^{g_0}(p) \setminus B_{c-b}^{g}(p)$,

(iii) $s(g_{d,s}^{b,\epsilon,c}) - s(g) \leq -s \cdot e^{-\frac{d}{a}}$ on $B_{c-a}^{g}(p) \setminus B_{c-b}^{g}(p)$,

Proof. (i) is clear from the definition. To show (ii) and (iii), one needs to consider terms in the right hand side of (10). From the assumption on g and Lemma 4 we have $|a_{ijk}| < \frac{2}{r_g}$ and $|a_{ijk,l}| < \frac{2}{r_g^2}$. Note that $0 \leq Y(r_g) \leq 1$ for $0 \leq s \leq 1$ and on $B_{10^4}^{g_0}(p)$, for any $d > 0$. So we have $|I_1| \leq \frac{\nu}{r_g^2}$ for a constant ν. By some computation ([12, Lemma 1.2 (i)]), there is $d_0(b) > 0$ such that for $d \geq d_0(b)$, $f_{d,s}^{(k)} > 0$ on $(0,b]$ for $k = 0,1,2,3$. We assert that there is $d_0(a,b,c) > d_0(b)$ such that for $d \geq d_0(a,b,c)$ and every $s > 0$ the following two inequalities hold:

$$f_{d,s}^{''} - f_{d,s}^{'}\frac{c_1}{c-t} - f_{d,s} \cdot \frac{\nu}{(c-t)^2} \geq 0, \quad \text{on } (0,b],$$

$$f_{d,s}^{''} - f_{d,s}^{'}\frac{c_1}{c-t} - f_{d,s} \cdot \frac{\nu}{(c-t)^2} \geq s \cdot e^{-\frac{d}{a}}, \quad \text{on } (a,b],$$

where c_1 is a constant such that $|3a_{122} + 2a_{133} + 2a_{144}| \leq \frac{c_1}{r_g}$. The proof of this assertion is similar to that of Lemma 1.2 (ii) in [12], so we omit it. Then we get on $B_c^g(p) \setminus B_{c-b}^g(p)$, where $h_\epsilon^b(c - r_g) \equiv 1$,

$$\begin{aligned} Y^{''} + Y^{'}(3a_{122}+&2a_{133} + 2a_{144}) + Y \cdot I_1 \\ &\geq f_{d,s}^{''}(c-r_g) - f_{d,s}^{'}(c-r_g)\frac{c_1}{r_g} - f_{d,s}(c-r_g) \cdot \frac{\nu}{(r_g)^2} \\ &= f_{d,s}^{''}(t) - f_{d,s}^{'}(t)\frac{c_1}{c-t} - f_{d,s}(t) \cdot \frac{\nu}{(c-t)^2}, \end{aligned}$$

where we set $t = c - r_g$. From above assertion we get (ii) and (iii). After all, we set $\gamma = \max\{1, d_0(a,b,c)\}$. ∎

5 Almost-Kähler island metrics on $\mathbb{R}^4$

In this section we describe almost Kähler island metrics on $\mathbb{R}^4$. We consider a metric on $\mathbb{R}^4$ of the form

$$\tilde{g} = f^2 dr^2 + \frac{r^2}{f^2} d\theta^2 + h^2 d\rho^2 + \frac{\rho^2}{h^2} d\sigma^2, \tag{11}$$

where $(r,\theta),(\rho,\sigma)$ are the polar coordinates for each summand of $\mathbb{R}^4 := \mathbb{R}^2 \times \mathbb{R}^2$ respectively and f, h are smooth positive functions on $\mathbb{R}^4$, which are functions of r and ρ only. Let $e_1 = \frac{1}{f}\frac{\partial}{\partial r}$, $e_2 = \frac{f}{r}\frac{\partial}{\partial \theta}$, $e_3 = \frac{1}{h}\frac{\partial}{\partial \rho}$, $e_4 = \frac{h}{\rho}\frac{\partial}{\partial \sigma}$. A smooth almost complex structure $J_{\tilde{g}}$ is defined by $J_{\tilde{g}}(e_1) = e_2, J_{\tilde{g}}(e_2) = -e_1, J_{\tilde{g}}(e_3) = e_4, J_{\tilde{g}}(e_4) = -e_3$. With the standard symplectic structure ω on $\mathbb{R}^4$ the triple $(\tilde{g}, \omega, J_{\tilde{g}})$ is an almost-Kähler structure on $\mathbb{R}^4$. Let ω_i be the dual co-frame field of e_i. Compute the connection 1-forms ω_{ij} w.r.t. ω_i: $d\omega_i = \sum_{j=1}^4 \omega_{ij} \wedge \omega_j$, with $\omega_{ij} = -\omega_{ji}$; one may compute $2a_{ijk} = \langle d\omega_k, \omega_i \wedge \omega_j\rangle_g - \langle d\omega_i, \omega_j \wedge \omega_k\rangle_g - \langle d\omega_j, \omega_k \wedge \omega_i\rangle_g$, where $\omega_{ij} = \sum_{k=1} a_{ijk}\omega_k$. And use the formula $d\omega_{ij} - \omega_{ik} \wedge \omega_{kj} = \sum_{k<l} R_{ijkl}\omega_k \wedge \omega_l$. Some Riemannian curvature components of $\tilde{g}$ are computed;

$$\begin{aligned}
R_{1212} &= -\frac{3f_r}{rf^3} + \frac{3f_r^2}{f^4} - \frac{f_{rr}}{f^3} - \frac{f_\rho^2}{f^2h^2}, \\
R_{1313} &= \frac{f_{\rho\rho}}{fh^2} - \frac{f_\rho h_\rho}{fh^3} + \frac{h_{rr}}{f^2h} - \frac{f_r h_r}{f^3h}, \\
R_{1414} &= -\frac{h_{rr}}{hf^2} + \frac{h_r f_r h + 2fh_r^2}{f^3h^2} + \frac{f_\rho}{\rho f h^2} - \frac{f_\rho h_\rho}{fh^3}, \\
R_{2424} &= -\frac{h_r}{rf^2h} + \frac{h_r f_r}{f^3h} - \frac{f_\rho}{\rho f h^2} + \frac{f_\rho h_\rho}{fh^3}, \\
R_{3434} &= -\frac{3h_\rho}{\rho h^3} - \frac{h_{\rho\rho}}{h^3} + \frac{3h_\rho^2}{h^4} - \frac{h_r^2}{f^2h^2}, \\
R_{2323} &= -\frac{f_{\rho\rho}}{fh^2} + \frac{f_\rho(2f_\rho h + fh_\rho)}{f^2h^3} + \frac{h_r}{rf^2h} - \frac{f_r h_r}{f^3h},
\end{aligned}$$

where $f_r = \frac{\partial f}{\partial r}, f_{rr} = \frac{\partial^2 f}{\partial r \partial r}$, etc.. The scalar curvature is then as follows;

$$s_{\tilde{g}} = -\{(f^{-2})_{rr} + \frac{3}{r}(f^{-2})_r + (h^{-2})_{\rho\rho} + \frac{3}{\rho}(h^{-2})_\rho\} - \frac{2f_\rho^2}{h^2f^2} - \frac{2h_r^2}{h^2f^2}.$$

Setting $F = f^{-2}$ and $H = h^{-2}$, we shall find F and H which satisfy

$$F_{rr} + \frac{3}{r} F_r + H_{\rho\rho} + \frac{3}{\rho} H_\rho = 0. \tag{12}$$

Set $F_{rr} + \frac{3}{r} F_r = \alpha(r)\beta(\rho)$ where α, β are smooth functions on $\mathbb{R}$ which satisfy at least that

$$\begin{aligned} \alpha(r) &= 0 \quad for \quad r \leq 0,\ r \geq 1, \\ \beta(\rho) &= 0 \quad for \quad \rho \leq 0,\ \rho \geq 1. \end{aligned}$$

These two functions α and β will be specified more below.

Since $(r^3 F_r)_r = r^3 F_{rr} + 3r^2 F_r = r^3 \alpha(r)\beta(\rho)$, we do integration with proper boundary conditions on F to get

$$F(r, \rho) = \beta(\rho) \int_0^r (\frac{1}{y^3} \int_0^y x^3 \alpha(x)\, dx)\, dy + 1.$$

We now specify the function α as follows; first consider a smooth function $p(y)$ on $\mathbb{R}$ such that

$$\begin{cases} a)\ p(y) = 0 \quad \text{for} \quad y \leq 0, y \geq 1, \\ b)\ |\frac{p'(y)}{y^3}| \ll 1,\ \text{for } y > 0 \\ c)\ \int_0^1 \frac{p(y)}{y^3}\, dy = 0, \\ d)\ 0 < \int_0^r \frac{p(y)}{y^3}\, dy < 1 \quad \text{for any} \quad r \quad \text{with} \quad 0 < r < 1 \end{cases}$$

and then define $\alpha(y) = \frac{p^{'}(y)}{y^3}$. Here we may choose one such function p so that the derivative $\alpha^{'}$ of α is zero at exactly three points in the open interval $(0, 1)$ as in Fig.1.



Fig.1. The graph of α.

Similarly we set

$$H(r,\rho) = -\alpha(r)\int_0^\rho (\frac{1}{y^3}\int_0^y x^3\beta(x)\,dx)\,dy + 1,$$

where $\beta(y) = \frac{k'(y)}{y^3}$ and the function $k(y)$ is a smooth function on $\mathbb{R}$ satisfying $a) - d)$. We may choose k similarly to p (or equally if one prefers) so that the derivative $\beta^{'}$ of β is zero at three points in $(0,1)$. Hence the graph of β is similar to that of α.

Then the functions $F(r,\rho)$ and $H(r,\rho)$ satisfy the equation (12) and

$$\begin{aligned} &F,\ H \equiv 1 \quad \text{for } r \leq 0,\ r \geq 1 \ \text{ or } \ \rho \leq 0,\ \rho \geq 1,\\ &F,\ H > 0 \quad \text{for } \ 0 < r < 1 \ \text{ and } \ 0 < \rho < 1. \end{aligned}$$

In particular, the metric $\tilde{g}$ is smooth on $\mathbb{R}^4$. For such functions F and H, we have

$$\begin{aligned} s_{\tilde{g}} = &-\frac{H}{2F^2}\beta'(\rho)^2\{\int_0^r (\frac{1}{y^3}\int_0^y x^3\alpha(x)\,dx)\,dy\}^2\\ &-\frac{F}{2H^2}\alpha'(r)^2\{\int_0^\rho (\frac{1}{y^3}\int_0^y x^3\beta(x)\,dx)\,dy\}^2. \end{aligned}$$

From the choices of α and β , the derivatives $\alpha^{'}(r)$ and $\beta^{'}(\rho)$ are zero at three points r_1, r_2, r_3 and ρ_1, ρ_2, ρ_3, respectively. It is now simple to check

$$s_{\tilde{g}}(r,\rho) = \begin{cases} 0 & \text{if } \mathrm{r} = 0,\ \mathrm{r} \geq 1,\ \rho = 0 \ \text{ or } \rho \geq 1,\\ 0 & \text{at } (\mathrm{r_i}, \rho_\mathrm{j}),\ \mathrm{i},\mathrm{j} = 1,2,3,\\ \text{negative} & \text{if } \ \mathrm{r}, \rho \in (0,1) \ \text{ with } \mathrm{r} \neq \mathrm{r_i} \text{ or } \rho \neq \rho_\mathrm{j}. \end{cases} \tag{13}$$

Now $\tilde{g}$ is an island metric and as such one may directly apply the main deformation of section 7 later, but to make our presentation clear we modify $\tilde{g}$ a little here. First, by choice of ingredient functions α and β, we can assume that $\tilde{g}$ satisfies $\|\tilde{g} - g_0\|_{C^{2\kappa+4}_{g_0}(\mathbb{R}^4)} \leq \frac{\lambda_3}{8}$, where g_0 is the Euclidean metric $dr^2 + r^2 d\theta^2 + d\rho^2 + \rho^2 d\sigma^2$.

We shall apply the deformation of section 4 to $\tilde{g}$. Choose a point p in $\mathbb{R}^4$ such that $r(p) = \rho(p) = \frac{r_1}{50}$. Then $s(\tilde{g})$ is negative on $B^{\tilde{g}}_{\frac{r_1}{100}}(p)$ by (4). In applying Lemma 5 and Lemma 6, we let $c = 1$, $a = 0.01$, $b = 1 - \frac{r_1}{200}$ and $\epsilon = \frac{r_1}{600}$, i.e. we consider the metric $\tilde{g}_{d,s}^{1-\frac{r_1}{200},\frac{r_1}{600},1}$ defined in terms of the $\tilde{g}$-orthonormal co-frame ω_i as defined in (5) with $\omega_1 = dr_{\tilde{g}}$ and $r_{\tilde{g}}(x) =$

$\mathrm{dist}_{\tilde g}(p,x)$. By Lemma 6, for $d = d_0 = \gamma(0.01, 1-\frac{r_1}{200}, 1)$ and $s \in [0,1]$ we have

(i) $\tilde g_{d,s}^{1-\frac{r_1}{200},\frac{r_1}{600},1} \equiv \tilde g$ on $B^{\tilde g}_{\frac{r_1}{300}}(p)$,

(ii) $s(\tilde g_{d,s}^{1-\frac{r_1}{200},\frac{r_1}{600},1}) - s(\tilde g) \leq 0$ on $\mathbb{R}^4 \setminus B^{\tilde g}_{\frac{r_1}{200}}(p)$,

(iii) $s(\tilde g_{d,s}^{1-\frac{r_1}{200},\frac{r_1}{600},1}) - s(\tilde g) \leq -s \cdot e^{-\frac{d}{a}}$ on $B^{\tilde g}_{0.99}(p) \setminus B^{\tilde g}_{\frac{r_1}{200}}(p)$.

From Lemma 5 (i) we have s_0 with $1 > s_0 > 0$ such that for $d = d_0 = \gamma(0.01, 1-\frac{r_1}{200}, 1), s^{-1} = s_0^{-1}$ we have

$$s(\tilde g_{d_0,s_0}^{1-\frac{r_1}{200},\frac{r_1}{600},1}) < 0, \text{ on } B^{\tilde g}_{\frac{r_1}{150}}(p) \quad \text{and,} \tag{14}$$

$$\|\tilde g_{d_0,s_0}^{1-\frac{r_1}{200},\frac{r_1}{600},1} - \tilde g\|_{C^{2\kappa+2}_{g_0}(\mathbb{R}^4)} < \frac{\lambda_3}{8}. \tag{15}$$

We shall denote by g^- the metric $\tilde g_{d_0,s_0}^{1-\frac{r_1}{200},\frac{r_1}{600},1}$. As $0 < r_1 < 1$, from above we deduce the following proposition;

Proposition 2 *With the standard symplectic structure on* $\mathbb{R}^4$ *there exists a compatible almost-Kähler metric* g^- *of non-positive scalar curvature on* $\mathbb{R}^4$ *which has the scalar curvature negative on the ball* $\{x|\ |x| < 0.9\}$ *and is Euclidean on* $\{x|\ |x| > 1.5\}$*. Furthermore, it satisfies*

$$\|g^- - g_0\|_{C^{2\kappa+2}_{g_0}(\mathbb{R}^4)} \leq \frac{\lambda_3}{4}. \tag{16}$$

We remark that the metric in (11) was first exhibited in [9]. Its scalar curvature, however, was not well presented, which should be as in above (13). So for completeness sake we rewrite it and apply the scalar-curvature diffusion, rather than supplying extra to the argument of that paper.

6 Almost Kähler Surgery

Given a compact almost Kähler manifold (M, g, ω, J) of real dimension 4, for each point $p \in M$ there exists an open neighborhood V_p in which there exists a coordinate system where ω has the standard form. Henceforth we shall call such a one a *Darboux* coordinate system. We choose an open coordinate ball of finite radius U_p such that $U_p \subset \overline{U_p} \subset V_p$, where $\overline{U_p}$ is the compact closure of U_p in M. These U_p's form a covering of M and there is a finite subcovering $\{U_{p_\alpha}\}_\alpha$, which we denote by $\mathfrak{A}$. We shall work with this (fixed) atlas. Denote the Lebesgue number of $\mathfrak{A}$ with respect to g by $\sigma := \sigma_{g,\mathfrak{A}}$.

Then, near each point p in M there is a Darboux coordinate system y_i in $\mathfrak{A}$ so that $\omega = \sum_{i=1}^{2} dy_{2i-1} \wedge dy_{2i}$ on $B^g_{\sigma/2}(p)$. Then g can be written as $g = \sum_{i,j=1}^{4} g_{ij} dy_i \otimes dy_j$. Consider the Euclidean metric $g_p = \sum_{i,j=1}^{4} g_{ij}(y(p)) dy_i \otimes dy_j$ on $B^g_{\sigma/2}(p)$, i.e. we extend $g|_p$, the restriction of g at p, onto $B^g_{\sigma/2}(p)$ with constant coefficients. Because y_i are Darboux coordinates, g_p is ω-almost Kähler with corresponding almost complex structure J_p, the extension of $J|_p$ onto $B^g_{\sigma/2}(p)$ with constant coefficients. Note that the definition of g_p depends on the choice of a Darboux coordinates system near p in $\mathfrak{A}$. However, there are only a finite number of coordinate systems in $\mathfrak{A}$, defined with the above property. Due to this aspect the following approximation Lemma holds, which provides the technical basis for subsequent Lemmas.

Lemma 7 *The following two statements hold.*

(i) For any $\varepsilon < 10^{-4}$, there is $m^{(0)} \geq \frac{10^5}{\sigma} > 0$ with $m^{(0)} = m^{(0)}(\varepsilon)$ such that for any $m > m^{(0)}$, $\|m^2 g - m^2 g_p\|_{C^{2\kappa+2}_{m^2 g_p}} < \varepsilon$ on $B^{m^2 g}_{10^4+1}(p)$ as well as on $B^{m^2 g_p}_{10^4}(p)$, where p is any point of M.

(ii) For any $\varepsilon < 10^{-4}$, there is $m^{(1)} \geq \frac{10^5}{\sigma} > 0$ with $m^{(1)} = m^{(1)}(\varepsilon)$ such that for any $m > m^{(1)}$, $\|m^2 g_q - m^2 g_p\|_{C^{2\kappa+2}_{m^2 g_p}} < \varepsilon$ on $B^{m^2 g}_{10^4+1}(p)$ as well as on $B^{m^2 g_p}_{10^4}(p)$, where p is any point of M and $q \in B^{m^2 g}_{2\cdot 10^4}(p)$.

Proof. Note that in the statement (ii) $m^2 g_q$ is defined on $B^{m^2 g}_{10^4+1}(p)$ because with $m \geq \frac{10^5}{\sigma}$, $m^2 g_q$ is defined on $B^{m^2 g}_{5\cdot 10^4}(q)$.

Let e_i, $i = 1, 2, 3, 4$ be a parallel orthonormal frame field of g_p. Then $e_i = \sum_{j=1}^{4} a_{ij} \frac{\partial}{\partial y_j}$, where a_{ij} are constants and y_j is any Darboux coordinates used to define g_p. By definition, in order to estimate $\|m^2 g - m^2 g_p\|_{C^0_{m^2 g_p}}$ on $B^{m^2 g}_{10^4+1}(p)$, we consider $| \, m^2 g(\frac{1}{m} e_i, \frac{1}{m} e_j) - m^2 g_p(\frac{1}{m} e_i, \frac{1}{m} e_j) \, | = | \sum_{s,t} a_{is} a_{jt} (g_{st} - g_{p\,st}) \, |$ on $B^g_{(10^4+1)/m}(p)$.

Note that $\max_{i,j} |a_{ij}|$ is determined by the continuous functions $g_{st} = g(\frac{\partial}{\partial y_s}, \frac{\partial}{\partial y_t})$ defined on a coordinate ball U_{p_α} of finite radius (there are only a finite number of such balls in $\mathfrak{A}$) and choice of an orthonormal frame. So there is a bound M, i.e. $|a_{ij}| \leq M$. For the same reason, for any $\varepsilon > 0$

there is $m^{(0)}$ such that $|\sum_{s,t} a_{is}a_{jt}(g_{st}-g_{p_{st}})| \leq M^2|\sum_{s,t}(g_{st}-g_{p_{st}})| \leq \varepsilon$ on $B^g_{(10^4+1)/m}(p)$ for $m > m^{(0)}$. The C^k-case ($k \geq 1$) can be done similarly. This proves (i) on $B^{m^2g}_{10^4+1}(p)$.

Next, we claim that $B^{m^2g_p}_{10^4}(p) \subset B^{m^2g}_{10^4+1}(p)$; consider the m^2g_p-geodesic $l(t)$ with $l(0)=p$ and $|l'(t)|_{m^2g_p} \equiv 1$. Let t_1 be the first value of t such that $l(t)$ lies in the boundary of $B^{m^2g}_{10^4+1}(p)$. Then

$$10^4+1 = d_{m^2g}(p, l(t_1)) \leq \int_0^{t_1} |l'(s)|_{m^2g} ds \leq \sqrt{1+\varepsilon}\int_0^{t_1} |l'(s)|_{m^2g_p} ds \leq \sqrt{1+\varepsilon}t_1.$$

As $10^4 < \frac{10^4+1}{\sqrt{1+\varepsilon}}$, this proves $B^{m^2g_p}_{10^4}(p) \subset B^{m^2g}_{10^4+1}(p)$, and finishes the proof of (i).

The proof of (ii) is similar, but note that g_p and g_q may use different Darboux coordinates, say y_i and z_i, respectively. We may write $g = \sum_{i,j} g_{ij}dy_i \otimes dy_j = \sum_{k,l} h_{kl}dz_k \otimes dz_l$, where $g_{ij} = \sum_{k,l} h_{kl}\frac{\partial z_k}{\partial y_i}\frac{\partial z_l}{\partial y_j}$. For the C^0-norm $\|m^2g_q - m^2g_p\|_{C^0_{m^2g_p}}$, it suffices to show that for m large $|m^2g_q(\frac{e_i}{m}, \frac{e_j}{m}) - m^2g_p(\frac{e_i}{m}, \frac{e_j}{m})| \leq \sum_{s,t} \mid a_{is}a_{jt} \mid \cdot \mid (g_{p_{st}} - g_{st}) + \sum_{k,l}(h_{kl} - h_{kl}(q))\frac{\partial z_k}{\partial y_s}\frac{\partial z_l}{\partial y_t} \mid$ is small on $B^g_{(10^4+1)/m}(p)$ for $q \in B^g_{2\cdot 10^4/m}(p)$. This can be done by a similar argument to (i). The C^k-case ($k \geq 1$) can be done similarly. ∎

For the given almost Kähler metric g on M, by Lemma 7 and Lemma 3 if $m \geq m^{(0)}(\lambda_1)$, the injectivity radius $\mathrm{inj_{m^2g}} \geq 200$ and the sectional curvature $|K_{m^2g}| \ll 1$.

We recall the Besicovitch covering Lemma 1. Assuming $m \geq m_0 \cdot m^{(0)}(\lambda_1)$, we shall explain a surgery replacing neighborhoods of $p \in A = A(m)$ in the almost Kähler manifold $(M, m^2g, m^2\omega, J)$ with copies of the metrics g^- of Proposition 2. Let $B_i := \{z \in A | B^{m^2g}_5(z) \in \mathcal{B}_i\}$. Then $A := \cup^{\kappa}_{i=1} B_i$.

At each point $p \in A$, there is a Darboux coordinate system y_i in $\mathfrak{A}$ such that on $B^{m^2g}_{10^4}(p)$, $m^2\omega = m^2\sum_i dy_{2i-1} \wedge dy_{2i}$, $m^2g = \sum_i m^2g_{ij}dy_idy^j$. And $m^2g_p = \sum_i m^2g_{ij}(p)dy_idy_j$ is a Euclidean metric compatible with $m^2\omega$ and the corresponding almost complex structure J_p. Now choose a g_p-parallel orthonormal frame field, e_1, $e_2 = J_pe_1$, e_3, $e_4 = J_pe_3$. There exists a coordinate system z_i such that $e_i = \frac{\partial}{\partial z_i}$. In this coordinate z_i, $m^2\omega = m^2\sum_i dz_{2i-1} \wedge dz_{2i}$ and $m^2g_p = \sum_i m^2dz_i \otimes dz_i$. Recall the metric g^- of Proposition 2 on $\mathbb{R}^4 = \{x = (x_1, x_2, x_3, x_4)\}$. Consider a diffeomorphism

$\phi = \phi_p : \{z|\ |z| < \frac{10^4}{m}\} \to \{x|\ |x| < 10^4\}$ defined by $z \mapsto mz$. Then from the definition of g^- we note that $\phi^* g^- = m^2 g_p$ outside $\{z||z| < \frac{1.5}{m}\}$.

From (2), we may express $m^2 g = (\phi^* g^-) \cdot e^h$ on $B^{m^2 g}_{10^4}(p)$ for a unique smooth symmetric $J_{\phi^* g^-}$-anti-invariant tensor h because $m^2 g$ and $\phi^* g^-$ are both $m^2\omega$-compatible. Let $\eta(r)$ be a smooth cutoff function in $C^\infty(\mathbb{R}^{>0}, [0,1])$ s.t. $\eta \equiv 0$ for $0 < r < 1.7$ and $\eta \equiv 1$ on the set $r \geq 1.8$.

For $m \geq m_0 \cdot m^{(0)}(\lambda_1)$ we define a metric on M;

$$g_A := \begin{cases} (\phi^* g^-) \mathrm{e}^{\eta(\mathrm{r}_{\mathrm{m^2 g}}) \cdot \mathrm{h}}, & \text{on } \cup_{\mathrm{p} \in \mathrm{A}} \mathrm{B}^{\mathrm{m^2 g}}_2(\mathrm{p}), \\ m^2 g, & \text{elsewhere.} \end{cases}$$

So we have embedded copies of the metric g^- into $(M, m^2 g, m^2\omega, J)$ near each $p \in A$.

Note that from Lemma 7 (i) and formula (4) we have $B^{m^2 g_p}_{1.5}(p) \subset B^{m^2 g}_{1.7}(p)$.

7 Main Deformation

Now we apply the scalar curvature diffusion of section 4, i.e. we deform g_A so that we decrease the scalar curvature away from neighborhoods of $p \in A$, and absorb the counter effect (scalar curvature increase) in neighborhoods of p where the scalar curvature of g_A is negative.

Using the Besicovitch covering $\mathcal{B}_1, \cdots, \mathcal{B}_\kappa$ we shall define ω-almost Kähler metrics $g(i)$ on M inductively on $i = 0, 1, 2, \cdots, \kappa$.

Observe that from (16) we have that for $m \geq m_0 \cdot m^{(0)}(\lambda_1)$,

$$\|\phi_q^* g^- - m^2 g_q\|_{C^{2\kappa+2}_{m^2 g_q}} \leq \frac{\lambda_3}{4} \qquad \text{on } B^{m^2 g_q}_{10^4}(q), \text{ for any } q \in A. \tag{17}$$

We set $g(0) = g_A$. We use often the following Lemma of base-metric change, whose proof is elementary.

Lemma 8 *Let g_1, g_2 be two Euclidean metrics on a domain Ω in $\mathbb{R}^4$ such that $B = B^{g_1}_{10^4}(p) \subset \overline{B} \subset \Omega$. For any $\delta > 0$ there exists $\varepsilon^{(0)} = \varepsilon^{(0)}(\delta)$ such that if $\|g_1 - g_2\|_{C^l_{g_1}(B)} < \varepsilon^{(0)}$, then $\|\eta\|_{C^l_{g_1}(\mathcal{O})} \leq (1+\delta)\|\eta\|_{C^l_{g_2}(\mathcal{O})}$ for any symmetric* (2,0)-tensor field η and any integer l with $2 \leq l \leq 2\kappa + 2$, where $\mathcal{O}$ is any open subset of B.

Now we shall make estimates on $g(0)$. Recall that on the region $B^{m^2g}_{1.8}(q)\setminus B^{m^2g}_{1.7}(q)$ the metric m^2g and the Euclidean metric $\phi_q^*g^- = m^2g_q$ are connected by a cutoff function. As $m^2g \to m^2g_q$ when $m\to\infty$, the next Lemma should be natural.

Lemma 9 *There exists* $\tilde{m}_0$ *with* $\tilde{m}_0 > m_0\cdot m^{(0)}(\lambda_1)$ *such that for* $m > \tilde{m}_0$, $\|g(0)-m^2g_q\|_{C^{2\kappa+2}_{m^2g_q}} \le \frac{\lambda_3}{4}$ *on* $B^{m^2g}_{1.8}(q)\setminus B^{m^2g}_{1.7}(q)$ *for each* $q\in A$.

Proof. On $U_q = B^{m^2g}_{1.8}(q)\setminus B^{m^2g}_{1.7}(q)$, we define h_m in $m^2g = (m^2g_q)e^{h_m}$. Then setting $m=1$, h_1 is defined by $g = g_qe^{h_1}$. One easily sees from definition (1) that $(m^2g_q)^{-1}h_m = (g_q)^{-1}h_1$ and $\frac{h_m}{m^2} = h_1$. Note that in general $(m^2g)e^h \neq m^2(ge^h)$.

If $\{e_i\}$ is a g_q-parallel orthonormal frame field, then $\{\frac{e_i}{m}\}$ is an m^2g_q-parallel orthonormal frame field. On $U_q = B^g_{\frac{1.8}{m}}(q)\setminus B^g_{\frac{1.7}{m}}(q)$, we have $m^2g(\frac{e_i}{m},\frac{e_j}{m}) = (m^2g_q)e^{h_m}(\frac{e_i}{m},\frac{e_j}{m}) = g_q(e_i, e^{(g_q)^{-1}h_1}(e_j))$ approach δ_{ij} as $m\to\infty$ uniformly, i.e. independently of point q and the coordinates chosen to define g_q. Or, simply it is Lemma 7 (i). Clearly $g_q \to g|_q$ and $(g_q)^{-1}h_1 \to 0$ uniformly.

Then, as $g(0)(\frac{e_i}{m},\frac{e_j}{m}) = (m^2g_q)\mathrm{e}^{\eta(\mathrm{r}_{\mathrm{m}^2\mathrm{g}})\cdot\mathrm{h}_\mathrm{m}}(\frac{\mathrm{e_i}}{\mathrm{m}},\frac{\mathrm{e_j}}{\mathrm{m}}) = \mathrm{g_q}(\mathrm{e_i}, \mathrm{e}^{\eta(\mathrm{r}_{\mathrm{m}^2\mathrm{g}})(\mathrm{g_q})^{-1}\mathrm{h_1}}(\mathrm{e_j}))$, so $g(0)$ approaches $m^2g_qe^0 = m^2g_q$ in the norm $\|\cdot\|_{C^0_{m^2g_q}}$. The proof of convergence in $\|\cdot\|_{C^{2\kappa+2}_{m^2g_q}}$ is similar and straightforward. ∎

The next Lemma is essential in the construction of the metrics $g(i)$'s;

Lemma 10 *There exists* m_1 *with* $m_1\ge\tilde{m}_0$ *such that for* $m\ge m_1$, $\|g(0) - m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}} \le \frac{\lambda_3}{2}$ *on* $B^{m^2g_p}_{10^4}(p)$ *for each* $p\in A$.

Proof. Fix $p\in A$. On $V_1 = B^{m^2g_p}_{10^4}(p)\setminus\cup_{q\in A}B^{m^2g}_{1.8}(q)$, $g(0) = m^2g$. By Lemma 7 (i), if $m\ge m^{(0)}(\frac{\lambda_3}{2})$ then $\|g(0)-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(V_1)} \le \frac{\lambda_3}{2}$.

On $B^{m^2g_p}_{10^4}(p)\cap B^{m^2g}_{1.7}(q)$ with $q\in A$, $g(0) = \phi_q^*g^-$, so we have the estimate (17) if $m\ge m_0\cdot m^{(0)}(\lambda_1)$. In the transition region $B^{m^2g_p}_{10^4}(p)\cap[B^{m^2g}_{1.8}(q)\setminus B^{m^2g}_{1.7}(q)]$, $q\in A$, Lemma 9 can be applied if $m>\tilde{m}_0$. So, on $W_q = B^{m^2g_p}_{10^4}(p)\cap B^{m^2g}_{1.8}(q)$ for $q\in A$ we estimated $g(0)-m^2g_q$ in terms of m^2g_q.

Now it remains to estimate $g(0)-m^2g_p$ on W_q with respect to m^2g_p. Applying Lemma 7 (ii) and Lemma 8 (with $g_1=m^2g_p$ and $g_2=m^2g_q$); if $m\ge m^{(1)}(\varepsilon^{(0)}(\frac{1}{2}))$;

$$\begin{aligned}\|g(0)-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(W_q)} &\le \|g(0)-m^2g_q\|_{C^{2\kappa+2}_{m^2g_p}(W_q)}+\|m^2g_q-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(W_q)}\\ &\le \frac{3}{2}\|g(0)-m^2g_q\|_{C^{2\kappa+2}_{m^2g_q}(W_q)}+\|m^2g_q-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(W_q)}\end{aligned}$$

By Lemma 7 (ii), $\|m^2g_q-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(W_q)}\le\frac{\lambda_3}{10\kappa}$ if $m\ge m^{(1)}(\frac{\lambda_3}{10\kappa})$.
From (17) and Lemma 9, for $m\ge m_1=\max\{\tilde{m}_0,\ m^{(1)}(\varepsilon^{(0)}(\frac{1}{2})),\ m^{(1)}(\frac{\lambda_3}{10\kappa})\}$ we have

$$\|g(0)-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(W_q)}\le\frac{3}{2}\cdot\frac{\lambda_3}{4}+\frac{\lambda_3}{10\kappa}<\frac{\lambda_3}{2}. \tag{18}$$

We proved that $\|g(0)-m^2g_p\|_{C^{2\kappa+2}_{m^2g_p}(B^{m^2g_p}_{10^4}(p))}\le\frac{\lambda_3}{2}$ for each $p\in A$ if $m\ge m_1$. $\blacksquare$

Suppose that for $m\ge m_1$, $g(j)$ is defined when $j=0,\cdots,i-1$ and satisfies the hypothesis

$$\|g(j)-m^2g_p\|_{C^{2\kappa+2-2j}_{m^2g_p}(B^{m^2g_p}_{10^4}(p))}\le\{\frac{1}{2}+\frac{j}{10\kappa}\}\lambda_3 \qquad \text{for } p\in A. \tag{19}$$

Lemma 10 corresponds to the case $j=0$. We note that in (19) $C^{2\kappa+2-2j}$-norm is used because in defining $g(1),\cdots,g(\kappa)$ successively we lose two derivatives at each step in applying Proposition 1 and Lemma 5, as we shall see in the proof of the next lemma.

Now, by Proposition 1 on each $B^{g(i-1)}_{100}(p)$ for $p\in B_i$, the $g(i-1)$-orthonormal co-frame field $\omega_1,\omega_2,\omega_3,\omega_4$ of (5) is defined. We set as in (8);

$$\begin{aligned}g(i)=&\frac{1}{(1+h^b_\epsilon\cdot f_{d,s}(9-d_{g(i-1)}(p,x)))}\omega_1\otimes\omega_1\\ &+(1+h^b_\epsilon\cdot f_{d,s}(9-d_{g(i-1)}(p,x)))\cdot\omega_2\otimes\omega_2 \quad +\omega_3\otimes\omega_3+\omega_4\otimes\omega_4.\end{aligned} \tag{20}$$

with $b=8.5, a=1, \epsilon=0.1, c=9$. With this done on $\cup_{p\in B_i}B^{g(i-1)}_{100}(p)$, $g(i)$ is smooth and ω-almost Kähler on M because $B^{g(i-1)}_9(p)$'s, $p\in B_i$ are disjoint by Lemma 1 (ii), Lemma 7 (i), the formula (4) and (19).

Lemma 11 *There exists $\tilde{d}$, independent of i, such that for $d \geq \tilde{d}$, $s \in (0,1]$ and $m \geq m_1$, the metric $g(i)$ satisfies the hypothesis* (19), i.e. (19) holds when $j = i$.

Proof. Consider $p \in A$ and let $z \in B_{10^4}^{m^2 g_p}(p)$. Set $V_z = B_{10^4}^{m^2 g_p}(p) \cap B_1^{m^2 g}(z)$.

Note that $V_z = V_z \cap (\cup_{j=1}^{\kappa} \cup_{q \in B_j} B_{10}^{m^2 g}(q)) = \cup_{j=1}^{\kappa} \{V_z \cap (\cup_{q \in B_j} B_{10}^{m^2 g}(q))\}$. $g(i) = g(i-1)$ away from $\cup_{q \in B_i} B_9^{g(i-1)}(q) \subset \cup_{q \in B_i} B_{10}^{m^2 g}(q)$. So, we get $\|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z)} = \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z \cap (\cup_{q \in B_i} B_{10}^{m^2 g}(q)))}$. Let Q be the set of q's in B_i such that $V_z \cap B_{10}^{m^2 g}(q) \neq \phi$. For $q \in Q$, from Lemma 8 and Lemma 7 (ii), if $m \geq m^{(1)}(\varepsilon^{(0)}(\frac{1}{2}))$,

$$\begin{aligned} \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(B_{10}^{m^2 g}(q))} &= \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(B_9^{g(i-1)}(q))} \\ &\leq \frac{3}{2} \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_q}(B_9^{g(i-1)}(q))}. \end{aligned}$$

And from the hypothesis (19) for $j = i-1$ and Lemma 5 (ii), there exists $\tilde{d} = d(\frac{2\lambda_3}{30\kappa}, 9)$ so that if $d \geq \tilde{d}$ and $s \in (0,1]$, $\|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_q}(B_9^{g(i-1)}(q))} \leq \frac{2\lambda_3}{30\kappa}$.
Combining above, we have

$$\begin{aligned} &\|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z \cap (\cup_{q \in B_i} B_{10}^{m^2 g}(q)))} \\ &\leq \sup_{q \in Q} \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(B_{10}^{m^2 g}(q))} \leq \frac{3}{2} \cdot \frac{2\lambda_3}{30\kappa} = \frac{\lambda_3}{10\kappa}. \end{aligned} \tag{21}$$

Finally, $\|g(i) - m^2 g_p\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z)}$

$$\begin{aligned} &\leq \|g(i) - g(i-1)\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z)} + \|g(i-1) - m^2 g_p\|_{C^{2\kappa+2-2i}_{m^2 g_p}(V_z)} \\ &\leq \frac{\lambda_3}{10\kappa} + \{\frac{1}{2} + \frac{i-1}{10\kappa}\}\lambda_3 = (\frac{1}{2} + \frac{i}{10\kappa})\lambda_3. \end{aligned}$$

So we proved (19) for $j = i$ and this proves the lemma. ∎

From Lemma 11, all the metrics $g(0)$, $g(1), \cdots, g(\kappa)$ can be defined. The parameters d and s are being used in defining each metric $g(i)$, $i = 1, \cdots, \kappa$. But we shall use the same values of d, s independently of i. Now we can prove;

Proposition 3 *There are m_2 with $m_2 > m_1$ and $d_1, s_1 > 0$ such that for each $d \geq d_1$, $s^{-1} \geq s_1^{-1}$ there exists an $m \geq m_2$ with $m = m(d,s)$ such that $s(g(\kappa)) < -c_1$ for a constant $c_1 = c_1(d,s) > 0$.*

Proof. For $\delta > 0$, there is an $m^{(2)} > m_1$ with $m^{(2)} = m^{(2)}(\delta)$ and

$$\sup_M |\ s(m^2 \cdot g)| < \delta \text{ and } -\mu - \delta < s(g(0)) < \delta, \tag{22}$$

for $m \geq m^{(2)}$ and $\mu := -\min\{\mathrm{s}(\mathrm{g}^-)\} > 0$. Here, for the second inequality one use the argument of Lemma 9, i.e. $g(0)$ approaches $m^2 g_q$ on the transition region $B_{1.8}^{m^2 g}(q) \setminus B_{1.7}^{m^2 g}(q)$ as m becomes large.

We consider $m > m_1$. Recall from Proposition 2 that $s(g^-) < 0$ on $\{x|\ |x| < 0.9\}$. So from Lemma 11, Lemma 5 (i) and Lemma 6 (ii) we have $\bar{d}_0$ with $\bar{d}_0 \geq \max\{\tilde{\mathrm{d}}, \gamma(1,\ 8.5,\ 9)\}$ and $\bar{s}_0$ with $1 \geq \bar{s}_0 > 0$ such that if we use d, s with $d \geq \bar{d}_0, s^{-1} \geq \bar{s}_0^{-1}$ in defining each metric $g(i)$, we have

$$s(g(\kappa)) < -\beta \ \text{ on } B_{0.8}^{m^2 g}(p),\ p \in A \tag{23}$$

where $\beta > 0$ is a constant.

By Lemma 6 for $d \geq \bar{d}_0$ and $s^{-1} \geq \bar{s}_0^{-1}$,

$$s(g(i+1)) - s(g(i)) \leq \begin{cases} -s \cdot e^{-d} \text{ on } \cup_{p \in B_{i+1}} B_8^{g(i)}(p) \setminus B_{0.5}^{g(i)}(p), \\ 0 \qquad\qquad\quad \text{on } M \setminus \cup_{p \in B_{i+1}} B_8^{g(i)}(p). \end{cases} \tag{24}$$

Note that the balls $B_8^{g(i)}(p)$, $p \in B_{i+1}$ are disjoint as $m > m_1$ from Lemma 1(ii), Lemma 7 (i), Lemma 11 and the formula (4). Now the rest of proof is almost the same as that in [12, proposition 4.5]. We produce it for completeness sake.

By Lemma 7 (i), Lemma 11 and the formula (4) again, we have $B_6^{m^2 g}(p) \setminus B_{0.55}^{m^2 g}(p) \subset B_8^{g(i)}(p) \setminus B_{0.5}^{g(i)}(p)$ for $p \in B^{i+1}$.

Therefore on $M \setminus \cup_{p \in A} B_{0.6}^{m^2 g}(p) \subset \cup_{p \in A} B_6^{m^2 g}(p) \setminus B_{0.55}^{m^2 g}(p)$, by adding inequalities (24), $i = 0, \cdots, \kappa - 1$, we get $s(g(\kappa)) - s(g(0)) \leq -s \cdot e^{-d}$. From (22) we have

$$s(g(\kappa)) < -s \cdot e^{-d} + \delta \quad \text{on } M \setminus \cup_{p \in A} B_{0.6}^{m^2 g}(p),$$

for any $\delta > 0$, $m \geq m^{(2)}(\delta)$ and $d \geq \bar{d}_0, s^{-1} \geq \bar{s}_0^{-1}$. Note that $\bar{d}_0, \bar{s}_0$ are independent of δ. Choose $\delta = \frac{s \cdot e^{-d}}{2}$. Then from (23) we obtain on M, $s(g(\kappa)) < -\min\{\frac{\mathrm{s} \cdot \mathrm{e}^{-\mathrm{d}}}{2}, \beta\}$.

Hence we will choose $d_1 = \bar{d}_0$, $s_1 = \bar{s}_0$, $m_2 = m^{(2)}(\delta_0)$ with $\delta_0 = \frac{\bar{s}_0 \cdot e^{-\bar{d}_0}}{2}$ and find that for each $d \geq d_1$, $s^{-1} \geq \bar{s}_1^{-1}$ there is an $m \geq m_2$ with

$$s(g(\kappa)) < -c_1 \qquad \text{for} \qquad c_1 = \min\{\frac{\mathrm{s} \cdot \mathrm{e}^{-\mathrm{d}}}{2}, \beta\}.$$

∎

We now have an almost Kähler structure $(g(\kappa), m^2\omega)$ of negative scalar curvature. The re-scaled metric $\frac{g(\kappa)}{m^2}$ is ω-compatible with negative scalar curvature. Theorem 1 is now proved.

Remark 1 Note that a Kähler version of the main theorem cannot hold. Indeed, we have the well known formula $\int_M s(g)\ dvol_g = 4\pi c_1(J) \cdot [\omega]$, where $c_1(J)$ is the first Chern class of a compact Kähler surface (M, g, ω, J) and $[\omega]$ is the cohomology class of ω. So, on the complex projective plane $\mathbb{CP}_2$ or a Hirzebruch surface F, a negative-scalar-curved Kähler metric does not exist, whereas negative-scalar-curved almost Kähler metrics exist on them by our main theorem.

Remark 2 In this article we only worked on the 4-dimensional case. And we wish to study the higher dimensional case in a forthcoming paper. It is interesting to pursue any implication of this result to symplectic topology.

Remark 3 From the nature of the scalar-curvature deformation argument of section 4, one may speculate that if there is a scalar-island metric in some category of metrics, then one may prove the existence of negative scalar curvature in the category. For instance one may study Hermitian metrics or contact metrics etc..

Remark 4 It is interesting to know if every closed symplectic manifold of dimension four admits a compatible almost Kähler metric of negative *Ricci* curvature or if the space of almost Kähler metrics with negative scalar curvature is contractible, as in [13].

Appendix

In this appendix we show the proof of Proposition 1 and supply details for the scalar curvature formula (9).

By Lemma 3, if we assume $\|g - g_0\|_{C^2_{g_0}(B^{g_0}_{10^4}(p))} \le \lambda_1$, then g has $\mathrm{inj}_g(p) \ge 200$. Let B denote the set $\{v \in T_p\mathbb{R}^4 \mid |v|_{g|_p} \le 100\}$, where $g|_p$ is the restriction of g to $T_p\mathbb{R}^4$. Then the g-exponential map at p, $\exp_p : B \to B^g_{100}(p) \subset \mathbb{R}^4$ has a smooth inverse $(\exp_p)^{-1}$. If we identify $T_p\mathbb{R}^4$ with $\mathbb{R}^4$ via the the map $i : T_p\mathbb{R}^4 \to \mathbb{R}^4$, $i(\sum_i v_i \frac{\partial}{\partial x_i}|_p) = (v_1, \cdots, v_4)$, then we can view $\exp_p$ as a map from a domain $i(B)$ in $\mathbb{R}^4$ to $\mathbb{R}^4$. For such a map we shall use the C^k norm $\|\cdot\|_{C^k}$ in the next lemma.

Lemma 12 *There exists* $\lambda^{(0)}$ *with* $\lambda^{(0)} < \lambda_1$ *and a positive constant* $C^{(0)}$ *such that if* $\|g - g_0\|_{C^l_{g_0}(B^{g_0}_{10^4}(p))} < \lambda \le \lambda^{(0)}$ *for* $4 \le l \le 2\kappa + 4$, *then* $\|\exp_p - (\mathrm{Id} + p)\ \|_{C^{l-1}(i(B))} < C^{(0)} \cdot \lambda$ *and* $\|(\exp_p)^{-1} - (\mathrm{Id} - p)\ \|_{C^{l-1}(B^g_{100}(p))} < C^{(0)} \cdot \lambda$. *Here,* $\mathrm{Id} + p$ *is the map* $(\mathrm{Id} + p)(x) = x + p$, *and likewise for* $\mathrm{Id} - p$.

Proof. To deal with the geodesic equation of the metric $g = \sum_{i,j=1}^4 g_{ij} dx_i \otimes dx_j$, i.e. $\ddot{x}_k = -\sum_{i,j=1}^4 \Gamma^k_{ij}(x(t))\dot{x}_i\dot{x}_j$, where Γ^k_{ij} is the Christoffel symbol of g, we consider the equivalent first order differential system:

$$\begin{cases} \dot{x}_k &= y_k \\ \dot{y}_k &= -\sum_{i,j} \Gamma^k_{ij}(x(t)) y_i y_j \end{cases} \tag{25}$$

with the initial condition $x(0) = p$ and $y(0) = v$. Denote its solution curve by $(x(t,v), y(t,v)) = (x_1(t,v), \cdots, x_4(t,v), y_1(t,v), \cdots, y_4(t,v))$.

Suppose that $\|g - g_0\|_{C^l_{g_0}} \le \varepsilon \le \lambda_1 < 10^{-4}$ with $4 \le l \le 2\kappa + 4$. Then, we have $|g_{ij} - \delta_{ij}| \le \varepsilon$, $|\frac{\partial^r g_{ij}}{\partial x_{m_1} \cdots \partial x_{m_r}}| \le \varepsilon$ for $r = 1, 2, \cdots, l$. For $\Gamma^k_{ij} = \frac{1}{2}\sum_{r=1}^4 g^{rk}(\frac{\partial g_{jr}}{\partial x_i} + \frac{\partial g_{ir}}{\partial x_j} - \frac{\partial g_{ij}}{\partial x_r})$, we have for $s = 0, 1, \cdots, l-1$,

$$|\frac{\partial^s \Gamma^k_{ij}}{\partial x_{m_1} \cdots \partial x_{m_s}}| \le c_1\varepsilon, \text{ with a positive constant } c_1. \tag{26}$$

In this appendix we shall use c_i, $i = 1, 2, \cdots$ to denote positive constants.

Since we study $\exp_p$ on B, it is enough to consider v with $|v| := |v|_{g_0} < 101$. And as $\exp_p(v) = x(1, v)$, we only need to consider t with $0 \le t \le 1$ when dealing with the solutions of (25). We write for convenience $\exp_p(v) = f(v) = (f_1(v), \cdots, f_4(v))$ and $\exp_p^{-1}(x) = h(x) = (h_1(x), \cdots, h_4(x))$.

Setting $Y(t) = |y(t)| = \sqrt{\sum_{i=1}^4 (y_i(t))^2}$, we get $|\dot{Y}| \le c_2\varepsilon\, Y^2$ from (25). Integrating it, we deduce from (25) $|\dot{y}_k| \le c_2\varepsilon Y^2 \le \frac{c_2\varepsilon|v|^2}{(1 - c_2\varepsilon t|v|)^2} \le c_3\varepsilon$, for small ε. So we get

$$\begin{aligned} &|y(t,v)-v| \le c_4\varepsilon, &&|x(t,v)-p-vt| \le c_4\varepsilon \\ &|f(v)-p-v| \le c_4\varepsilon, &&|h(x)-(x-p)| \le c_5\varepsilon. \end{aligned} \tag{27}$$

where the last inequality is deduced from the others and $h \circ f(v) = v$.

Now, the solution $x(t,v)$ and $y(t,v)$ of (25) are well known to be C^∞ functions of both variables t and v. Let p_k and v_k be the k-th component of p and v, respectively. From (25) we get for any n with $1 \le n \le l-1$,

$$\begin{cases} \frac{\partial^n x_k}{\partial v_{i_1}\cdots\partial v_{i_n}} &= \int_0^t \frac{\partial^n y_k}{\partial v_{i_1}\cdots\partial v_{i_n}}\, ds \\ \frac{\partial^n (y_k - v_k)}{\partial v_{i_1}\cdots\partial v_{i_n}} &= \int_0^t \sum_{m,i,j} -\frac{\partial \Gamma^k_{ij}}{\partial x^m}(\frac{\partial^n x_m}{\partial v_{i_1}\cdots\partial v_{i_n}}) y_i y_j - 2\Gamma^k_{ij}(\frac{\partial^n y_i}{\partial v_{i_1}\cdots\partial v_{i_n}}) y_j + L_n ds \end{cases} \tag{28}$$

where L_n consists of terms involving partial derivatives of x and y of order lower than n only. Note that $L_1 = 0$. We shall prove the following for $|v| < 101, 0 \le t \le 1$ when $0 \le m \le l-1$;

$$|\frac{\partial^m (y_k - v_k)}{\partial v_{i_1}\cdots\partial v_{i_m}}| \le \text{constant}\cdot\varepsilon, \qquad |\frac{\partial^m (x_k - p_k - v_k t)}{\partial v_{i_1}\cdots\partial v_{i_m}}| \le \text{constant}\cdot\varepsilon. \tag{29}$$

We will do this by induction on $m \ge 0$; The case $m = 0$ holds by (27). Suppose that (29) holds for $m = n-1 \le l-2$, for some integer $n \ge 1$. Then $|\int_0^t L_n ds| \le c_6\varepsilon$ from induction hypothesis. From the first equation in (28), $|\frac{\partial^n x_k}{\partial v_{i_1}\cdots\partial v_{i_n}}(t,v)| \le \sup_{s\in[0,t]} |\frac{\partial^n y_k}{\partial v_{i_1}\cdots\partial v_{i_n}}(s,v)|$, as $0 \le t \le 1$. So, using (27), from the second equation in (28) we deduce;

$$\sum_{k=1}^4 \sup_{s\in[0,t]} |\frac{\partial^n (y_k - v_k)}{\partial v_{i_1}\cdots\partial v_{i_n}}(s,v)| \le c_7\varepsilon \sum_{k=1}^4 \sup_{s\in[0,t]} |\frac{\partial^n y_k}{\partial v_{i_1}\cdots\partial v_{i_n}}(s,v)| + 4c_6\varepsilon. \tag{30}$$

From this, for small ε we easily derive (29) when $m = n$. Then, for any $0 \le m \le l-1$,

$$|\frac{\partial^m (f_k(v) - (p_k + v_k))}{\partial v_{i_1}\cdots\partial v_{i_m}}| \le c_8\varepsilon \quad \text{and} \quad |\frac{\partial^m (h_k(x) - (x_k - p_k))}{\partial x_{j_1}\cdots\partial x_{j_m}}| \le c_9\varepsilon. \tag{31}$$

where the second inequality is deduced from the first, using (differentiation of) $h \circ f(v) = v$. We let $\lambda^{(0)}$ to be a small ε value which makes all above arguments hold. Choice of $C^{(0)}$ should be obvious. This proves the lemma. ■

Lemma 13 *There exist* $\lambda^{(1)}$ *with* $\lambda^{(1)} < \lambda_1$ *and a positive constant* $C^{(1)}$ *such that if* $\|g - g_0\|_{C^l_{g_0}} \leq \lambda \leq \lambda^{(1)}$ *on* $B^{g_0}_{10^4}(p) \subset \mathbb{R}^n$*, where* $4 \leq l \leq 2\kappa + 4$*, then for* $j = 0, 1, \cdots, l-2$*,*

$$\|(\nabla^{g_0})^j dr_g - (\nabla^{g_0})^j dr_{g_0}\|_{C^0_{g_0}} < \frac{C^{(1)} \cdot \lambda}{r^j_{g_0}} \qquad \text{on } B^g_{100}(p) - \{p\}. \tag{32}$$

Proof. We choose $\lambda^{(1)}$ to be $\lambda^{(0)}$ of Lemma 12. We have $r_g(x) = | \exp_p^{-1}(x)|_{g|_p} = \sqrt{\sum_{i,j=1}^4 g_{ij}(p)\ h_i(x)h_j(x)}$ for x with $r_g(x) < 200$. We shall prove (32) by induction on j. The case $j = 0$ can be checked easily by applying (31). Suppose that (32) holds for $j = 0, \cdots, i-1 \leq l-3$. For the rest of this proof we write $\partial^k f$ to denote a k-th order partial derivative of a function f in x coordinates, by an abuse of terminology. Then we may write for $\mu = g$ or g_0

$$\partial^{i+1}(r^2_\mu) - 2r_\mu(\partial^{i+1} r_\mu) = \sum_{k=1}^{i} a_k (\partial^k r_\mu)(\partial^{i+1-k} r_\mu) \tag{33}$$

where a_k's are natural numbers. Here ∂^{i+1} in the LHS is a specific partial derivative, e.g. $\frac{\partial^{i+1} r^2_\mu}{\partial x_{s_1} \cdots \partial x_{s_{i+1}}}$ but the index k in the RHS should be understood as a multi-index. We have

$$\partial^{i+1} r_g - \partial^{i+1} r_{g_0} - \frac{\partial^{i+1}({r_g}^2)}{2r_g} + \frac{\partial^{i+1}(r^2_{g_0})}{2r_{g_0}} = \sum_{k=1}^{i} \frac{a_k}{2} \left(\frac{\partial^k r_{g_0} \partial^{i+1-k} r_{g_0}}{r_{g_0}} - \frac{\partial^k r_g \partial^{i+1-k} r_g}{r_g}\right). \tag{34}$$

Using (33) for $\mu = g_0$ and $\partial^{i+1} r^2_{g_0} = 0$ for $i \geq 2$, one can derive, by induction;

$$|\partial^{i+1} r_{g_0}| \leq \frac{c_{10}}{r^i_{g_0}} \text{ for } i = 0, 1, \cdots, l-2. \tag{35}$$

Using this inequality and the induction hypothesis, $\sum_{k=1}^i a_k |\frac{\partial^k r_{g_0} \partial^{i+1-k} r_{g_0}}{r_{g_0}} - \frac{\partial^k r_g \partial^{i+1-k} r_g}{r_g}| \leq \frac{c_{11}\lambda}{r^i_{g_0}}$. We need to show $|\frac{\partial^{i+1}(r^2_g)}{r_g} - \frac{\partial^{i+1}(r^2_{g_0})}{r_{g_0}}| \leq \frac{\text{constant}\cdot\lambda}{r^i_{g_0}}$. One can check easily from (31) that $r^2_g = \sum_{i,j=1}^4 g_{ij}(p)\ h_i(x)h_j(x)$ satisfies $|\frac{\partial^2(r^2_g)}{r_g} - \frac{\partial^2(r^2_{g_0})}{r_{g_0}}| \leq \frac{c_{12}\lambda}{r_{g_0}}$. For $i \geq 2$, $|\partial^{i+1}(r^2_g)| \leq c_{13}\lambda$ from (31). As $\partial^{i+1}(r^2_{g_0}) = 0$, we have $|\frac{\partial^{i+1}(r^2_g)}{r_g} - \frac{\partial^{i+1}(r^2_{g_0})}{r_{g_0}}| = |\frac{\partial^{i+1}(r^2_g)}{r_g}| \leq \frac{c_{13}\lambda}{r_g} \leq \frac{c_{14}\lambda}{r^i_{g_0}}$ on $B^g_{100}(p) - \{p\}$. In sum, from (34) we get $|\partial^{i+1} r_g - \partial^{i+1} r_{g_0}| \leq \frac{c_{15}\lambda}{r^i_{g_0}}$. This proves (32) for some $C^{(1)}$

when $0 \le j \le l-2 \le 2\kappa + 2$ and the Lemma. ∎

Proof of Proposition 1; In Lemma 13 we already proved the inequality (6) for ω_1. From the hypothesis on g we have $\|J_g - J_0\|_{C^l_{g_0}} \le \lambda$, so the proposition holds for $\omega_2 = J_g\omega_1$. With these results for ω_1 and ω_2, by choosing λ_2 small enough, the denominator of ω_3 is never zero on $B^g_{100}(p) - \{p\}$. It is now a straightforward computation to show (6) for ω_3 and ω_4. This finishes the proof of Proposition 1.

Now we shall derive the formula (9) in section 4. For the metric $g = \sum_{i=1}^4 \omega_i \otimes \omega_i$ in section 4 we write $d\omega_i = \sum_j \omega_{ij} \wedge \omega_j$ with $\omega_{ij} = -\omega_{ji}$ and similarly for $\tilde g := g^{b,\epsilon,c}_{d,s}$, $d\tilde\omega_i = \sum_j \tilde\omega_{ij} \wedge \tilde\omega_j$ with $\tilde\omega_{ij} = -\tilde\omega_{ji}$. We set $\omega_{ij} = \sum_k a_{ijk}\omega_k$ and $\tilde\omega_{ij} = \sum_k \tilde a_{ijk}\tilde\omega_k$. Then,

$$d\omega_i = \sum_{j<k}(a_{ikj} - a_{ijk})\omega_j \wedge \omega_k \quad \text{and} \quad d\tilde\omega_i = \sum_{j<k}(\tilde a_{ikj} - \tilde a_{ijk})\tilde\omega_j \wedge \tilde\omega_k. \tag{36}$$

From (5), $d\omega_1 = d(dr_g) = \sum_{j<k}(a_{1kj} - a_{1jk})\omega_j \wedge \omega_k = 0$, so $a_{1jk} = a_{1kj}$ for any j,k. We let $\alpha(r_g) = \frac{1}{\sqrt{1+Y(r_g)}} = \frac{1}{\sqrt{1+h^b_\epsilon \cdot f_{d,s}(c-r_g)}}$ so that $\tilde\omega_1 = \alpha(r_g)\omega_1$. Then $d\tilde\omega_1 = d(\alpha(r_g)dr_g) = 0$, so $\tilde a_{1jk} = \tilde a_{1kj}$. Comparing the two equations in (36), one gets the following relations. Other ones can be derived from these, using $a_{ijk} = -a_{jik}$ and $a_{1jk} = a_{1kj}$.

$$\text{For } i = 2,3,4, \quad \tilde a_{1i1} = 0, \qquad a_{1i1} = 0.$$

$$\text{For } j = 3,4, \qquad \tilde a_{2j2} = a_{2j2}, \quad \tilde a_{j1j} = \frac{1}{\alpha}a_{j1j}, \quad \tilde a_{j2j} = \alpha a_{j2j},$$

$$\tilde a_{2j1} = \frac{1}{2}(1 - \frac{1}{\alpha^2})a_{21j} + \frac{1}{2}(1 + \frac{1}{\alpha^2})a_{2j1}, \quad \tilde a_{21j} = \frac{1}{2}(1 + \frac{1}{\alpha^2})a_{21j} + \frac{1}{2}(1 - \frac{1}{\alpha^2})a_{2j1}.$$

$$\tilde a_{212} = \frac{\alpha'}{\alpha^2} + \frac{a_{212}}{\alpha}, \qquad \tilde a_{343} = a_{343}, \quad \tilde a_{434} = a_{434}, \quad \tilde a_{143} = \frac{1}{\alpha}a_{143}, \quad \tilde a_{341} = \frac{1}{\alpha}a_{341}$$

$$\tilde a_{342} = \frac{1}{2\alpha}(a_{243} - a_{234}) + \frac{\alpha}{2}(2a_{342} + a_{234} - a_{243}),$$

$$\tilde a_{234} = \frac{1}{2\alpha}(a_{234} - a_{243}) + \frac{\alpha}{2}(a_{243} + a_{234}),$$

$$\tilde a_{243} = \frac{1}{2\alpha}(a_{243} - a_{234}) + \frac{\alpha}{2}(a_{243} + a_{234}).$$

We compute the curvature components $\tilde R_{ijij}$ of $g^{b,\epsilon,c}_{d,s}$ from $d\tilde\omega_{ij} - \sum_k \tilde\omega_{ik} \wedge \tilde\omega_{kj} =$

$\sum_{k<l} \tilde{R}_{ijkl}\tilde{\omega}_k \wedge \tilde{\omega}_l$ and the scalar curvatue. We set $d(\tilde{a}_{ijk}) = \sum_s \tilde{a}_{ijk,s}\tilde{\omega}_s$.

$$\begin{aligned}
\tilde{R}_{1212} =& \tilde{a}_{122,1} + \tilde{a}_{122}^2 + \tilde{a}_{123}^2 + \tilde{a}_{124}^2 + 2(\tilde{a}_{123}\tilde{a}_{321} + \tilde{a}_{124}\tilde{a}_{421}),\\
\tilde{R}_{1313} =& \tilde{a}_{133,1} + \tilde{a}_{132}^2 + \tilde{a}_{133}^2 + \tilde{a}_{134}^2 + 2(\tilde{a}_{132}\tilde{a}_{231} + \tilde{a}_{134}\tilde{a}_{431}),\\
\tilde{R}_{1414} =& \tilde{a}_{144,1} + \tilde{a}_{142}^2 + \tilde{a}_{143}^2 + \tilde{a}_{144}^2 + 2(\tilde{a}_{142}\tilde{a}_{241} + \tilde{a}_{143}\tilde{a}_{341}),\\
\tilde{R}_{2323} =& \tilde{a}_{233,2} - \tilde{a}_{232,3} + \tilde{a}_{234}(\tilde{a}_{432} - \tilde{a}_{423}) + \tilde{a}_{232}^2 + \tilde{a}_{233}^2 + \tilde{a}_{122}\tilde{a}_{133} + \tilde{a}_{422}\tilde{a}_{433}\\
&- \tilde{a}_{123}^2 + \tilde{a}_{243}\tilde{a}_{432},\\
\tilde{R}_{2424} =& \tilde{a}_{244,2} - \tilde{a}_{242,4} + \tilde{a}_{243}(\tilde{a}_{342} - \tilde{a}_{324}) + \tilde{a}_{242}^2 + \tilde{a}_{244}^2 + \tilde{a}_{122}\tilde{a}_{144} + \tilde{a}_{322}\tilde{a}_{344}\\
&- \tilde{a}_{124}^2 + \tilde{a}_{234}\tilde{a}_{342},\\
\tilde{R}_{3434} =& \tilde{a}_{344,3} - \tilde{a}_{343,4} + \tilde{a}_{342}(\tilde{a}_{243} - \tilde{a}_{234}) + \tilde{a}_{343}^2 + \tilde{a}_{344}^2 + \tilde{a}_{133}\tilde{a}_{144} + \tilde{a}_{233}\tilde{a}_{244}\\
&- \tilde{a}_{134}^2 + \tilde{a}_{324}\tilde{a}_{243},
\end{aligned}$$

$$\begin{aligned}
\frac{s(g_{d,s}^{b,\epsilon,c})}{2} =& \tilde{R}_{2112} + \tilde{R}_{3113} + \tilde{R}_{4114} + \tilde{R}_{3223} + \tilde{R}_{4224} + \tilde{R}_{4334}\\
=& -\sum_{i=2}^{4} \tilde{a}_{1ii,1} - \sum_{2\le i\le j\le 4} \tilde{a}_{1ij}^2 + \sum_{i=3}^{4}(\tilde{a}_{2i2,i} - \tilde{a}_{2ii,2}) + \tilde{a}_{343,4} - \tilde{a}_{344,3}\\
&+ \tilde{a}_{423}\tilde{a}_{342} + \tilde{a}_{234}\tilde{a}_{342} + \tilde{a}_{234}\tilde{a}_{423} - \sum_{i=3}^{4}(\tilde{a}_{2i2}^2 + \tilde{a}_{2ii}^2) - \tilde{a}_{343}^2 - \tilde{a}_{344}^2\\
&- \sum_{i=3}^{4} \tilde{a}_{122}\tilde{a}_{1ii} - \tilde{a}_{422}\tilde{a}_{433} - \tilde{a}_{322}\tilde{a}_{344} - \tilde{a}_{133}\tilde{a}_{144} - \tilde{a}_{233}\tilde{a}_{244}\\
=& \frac{\alpha^{''}}{\alpha^3} - 3\frac{(\alpha^{'})^2}{\alpha^4} + \frac{\alpha^{'}}{\alpha^3}(3a_{122} + 2a_{133} + 2a_{144}) - \frac{1}{\alpha^2}\sum_{i=2}^{4}(a_{1ii,1} + a_{1ii}^2)\\
&- \sum_{i=3}^{4} \frac{1}{4}\{(1 + \frac{1}{\alpha^2})a_{21i} + (1 - \frac{1}{\alpha^2})a_{2i1}\}^2 - \frac{1}{\alpha^2}a_{134}^2\\
&+ \sum_{i=3}^{4}(a_{2i2,i} - \alpha^2 a_{2ii,2}) + a_{343,4} + a_{434,3} + (\frac{1}{2} - \frac{1}{4\alpha^2})(a_{243} - a_{234})^2\\
&+ a_{342}(a_{234} - a_{243}) - \frac{\alpha^2}{4}(a_{243} + a_{234})^2 - \sum_{i=3}^{4}(\alpha^2 a_{2ii}^2 + a_{2i2}^2) - a_{343}^2\\
&- a_{344}^2 - \sum_{i=3}^{4} \frac{a_{122}}{\alpha^2}a_{1ii} + a_{242}a_{433} + a_{232}a_{344} - \frac{1}{\alpha^2}a_{133}a_{144} - \alpha^2 a_{233}a_{244}.
\end{aligned}$$

Putting $\frac{1}{\alpha^2} = 1 + Y$, $\frac{\alpha'}{\alpha^3} = -\frac{1}{2}Y'$ and $\frac{\alpha''}{\alpha^3} - 3\frac{(\alpha')^2}{\alpha^4} = -\frac{1}{2}Y''$ into the above, we get the formula (9).

References

[1] V. Apostolov, T. Drăghici, D. Kotschick, *An integrability theorem for almost Kahler* 4*-manifolds.* C. R. Acad. Sci. Paris Ser. I Math. 329 (1999), no. 5, 413–418.

[2] A.L. Besse, Einstein manifolds, Ergebnisse der Mathematik, 3. Folge, Band 10, Springer-Verlag, 1987.

[3] D.E. Blair, *On the set of metrics associated to a symplectic or contact form,* Bull. Institute of Math. Academia Sinica, Vol.11, No.3, Sep. (1983), 297-308.

[4] J. Cheeger, D. G. Ebin, Comparison Theorems in Riemannian Geometry, North-Holland Publ., Volume 9, Springer-Verlag, 1975.

[5] J. Cheeger, M. Gromov, and M. Taylor, *Finite propagation speed, kernel estimates for functions of the Laplace operator and the geometry of complete Riemannian manifolds*, J. Differential Geom. 17 (1982), 15-53.

[6] R.E. Gompf and A.I. Stipsicz, *4-Manifolds and Kirby Calculus*, Graduate Studies in Math., v.20, American Math. Soc., 1999.

[7] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. 82 (1985), 307-347.

[8] M. Itoh, *Almost Kahler 4-manifolds,* L^2*-scalar curvature functional and Seiberg-Witten equations.* Internat. J. Math. 15 (2004), no. 6, 573–580.

[9] Y. Kang and J. Kim *Almost Kahler metrics with non-positive scalar curvature which are Euclidean away from a compact set*, Jour. Korean. Math. Soc. 41 no.5, (2004), 809-820

[10] J. Kim, C. Sung, *Deformations of almost-Kahler metrics with constant scalar curvature on compact Kahler manifolds.* Ann. Global Anal. Geom. 22 (2002), no. 1, 49–73.

[11] W. Klingenberg, *Contributions to Riemannian geometry in the large*, Ann. of Math. (2) 69 (1959), 654-666

[12] J. Lohkamp, *Metrics of negative Ricci curvature*, Ann. of Math. (2) 140 (1994), 655-683.

[13] J. Lohkamp, *The space of negative scalar curvature metrics*, Invent. Math. 110 (1992), no. 2, 403-407.

[14] K. Yano, *Differential Geometry on Complex and Almost Complex Spaces*, Pergamon Press Ltd, 1965